

Dry Bean N Fertility



on-farm network
PARTICIPATORY • PRECISE • PROACTIVE

***What's the optimum N rate for dry bean production?
It might be lower than you think. Find out!***

Suggested Treatments

- 1) Normal rate of N
- 2) Low N rate: normal - 30 lb N/ac
- 3) High N rate: normal + 30 lb N/ac

Trial Design

- Follow a randomized layout
- Minimum four replicates
- Strips need to be at least as wide as one combine pass
- Strips should run the length of the field, excluding the headlands

Headland

Normal Rate	Low Rate	High Rate	Low Rate	High Rate	Normal Rate	Low Rate	Normal Rate	High Rate	Normal Rate	Low Rate	High Rate
1	2	3	4	5	6	7	8	9	10	11	12

Headland

***Interested in different rates?
Let us know!***

Your Fields, Your Results

- Established with your equipment
- Under your management practices
- Harvested with your equipment
- MPSG support at establishment & harvest
- MPSG collects data throughout the season
- Results packaged together for you in a short report
- Must be a member in good standing

***Decision Making, Simplified
Your trial includes:***

- ✓ Plant stand data
- ✓ Nodulation ratings
- ✓ Yield results
- ✓ Soil testing
- ✓ Economics analysis

Your data, working for your farm

MANITOBA
Pulse Soybean
GROWERS

To participate in the trials, contact

Leanne Koroscil • 204.751.0439 • leanne@manitobapulse.ca

or sign up at manitobapulse.ca/on-farm-network